arXiv:1512.00323v3 [math.AG] 9 May 2016

THE DUAL COMPLEX OF $\overline{M}_{0,n}$ VIA PHYLOGENETICS

NOAH GIANSIRACUSA

ABSTRACT. The moduli space $\overline{M}_{0,n}$ of stable rational n-pointed curves has divisorial boundary with simple normal crossings. In this brief note I observe that the dual complex is a flag complex; that is, a collection of boundary divisors has nonempty intersection if and only if the pairwise intersections are nonempty. Rather than proving this directly, I translate the statement to a setting in phylogenetics where it is widely used and multiple explicit proofs have been written. It appears this result is known by experts but lacks a detailed reference in the literature, except recently for $n = 7$.

1. INTRODUCTION

The moduli space $\overline{M}_{0,n}$ of stable rational n-pointed curves has featured in many areas of mathematics. It was first envisioned by Grothendieck [GRR72], then constructed as an iterated blow-up by Knudsen [Knu83], then given slicker iterated blow-up constructions by Kapranov [Kap93a, Kap93b] and Keel [Kee92]. It has been the subject of interest in birational geometry [KM13, CT15], arithmetic [GM04], topology [EHKR10], representation theory [AFV11], enumerative geometry [KV07], derived categories [MS13], etc. Its scope even reaches beyond pure mathematics, as the tree structures arising in $\overline{M}_{0,n}$ play an important role in phylogenetics [BHV01, Kap93c, SS04, LP11]. In this brief note I exploit this latter connection to provide a succinct proof, via reference to a combinatorial result in the phylogenetics literature, of a folklore result about $\overline{M}_{0,n}$: a collection of boundary divisors has nonempty intersection if and only if all pairs in the collection have nonempty intersection. This can be stated formally as follows (see [Kol14] for a nice discussion of dual complexes and their applications and [KV07] for an elementary introduction to $\overline{M}_{0,n}$, and recall that a simplicial complex is called a flag complex if it is the maximal simplicial complex on its 1-skeleton):

Theorem. *The dual complex of the divisorial boundary in $\overline{M}_{0,n}$ is a flag complex.*

The really is a special property; for instance, the three coordinate lines in $\mathbb{P}^2$ intersect in pairs whereas their triple intersection is empty. This fact about $\overline{M}_{0,n}$ seems to be known by some experts (cf., [GM10, p.871], [HKT09, Lemma 7.2], [SS04]) and was recently proven explicitly in the first non-trivial case, $n = 7$ [Sch15, Lemma 3.2 and Proposition 3.4]. As I show below, by considering the well-known stratification of $\overline{M}_{0,n}$ by boundary strata corresponding to dual graph trees, this intersection-theoretic property of $\overline{M}_{0,n}$ is precisely the combinatorial "Splits-Equivalence Theorem" from the 70s due to Buneman [Bun71, Bun74], which in turn is the "Four Gamete Condition" in phylogenetics [EJM76, Gus91, Mea83]. Buneman's result was given an alternate proof in the 80s [BD86] which is nicely exposed in Semple and Steel's book on mathematical phylogenetics [SS03].

Acknowledgements. I thank Luca Schaffler for bringing to my attention this question about $\overline{M}_{0,n}$ and I thank Luca, Jenia Tevelev, and James McKernan for helpful discussions. I also thank the Simons Foundation for financial support.

MSC-class: 14H10, 05C05

2. Proof via reinterpretation

A *phylogenetic tree on n taxa* is a graph that is a tree with n leaves, labelled by the integers $1,\dots,n$, such that there are no vertices of degree two. Each point of $\overline{M}_{0,n}$ determines a phylogenetic tree [Kap93c, §4]. Indeed, the points of $\overline{M}_{0,n}$ are in bijection with stable rational n-pointed curves $(C,p_1,\dots,p_n)$, which means C is a nodal union of $\mathbb{P}^1$s such that each irreducible component contains at least three *special* points (nodes or marked points p_i) and such that the dual graph is a tree. The *dual graph* here refers to the graph with a vertex for each irreducible component $C_i \subseteq C$ and each marked point p_i, and an edge joining two vertices if they correspond to irreducible components satisfying $C_i \cap C_j \neq \emptyset$ or to a component and a marked point satisfying $p_j \in C_i$. Such an object is necessarily a phylogenetic tree. These dual graphs induce a widely-used stratification of $\overline{M}_{0,n}$, where a *stratum* is the closure of the locus of points with fixed dual graph. The strata in $\overline{M}_{0,n}$ are in bijection with the phylogenetic trees on n taxa, where the bijection sends a stratum to the dual graph of any point in its relative interior (which we shall refer to simply as the dual graph of the stratum).

In the phylogenetics literature, a bipartition $I \sqcup J = \{1,\dots,n\}$ is referred to as a *split*. Each *internal* edge (that is, edge between non-leaf vertices) of a phylogenetic tree induces a split by considering the sets of integer labels on either side of the edge (which is well-defined due to the tree property of the graph). The set of all such splits of a phylogenetic tree T is denoted $\Sigma(T)$. A pair of splits $I \sqcup J$, $K \sqcup L$ is said to be *compatible* if at least one of the following intersections is empty (see [SS03, Chapter 3]): $I \cap K$, $I \cap L$, $J \cap K$, $J \cap L$. The famous Buneman Splits-Equivalence Theorem alluded to in the introduction is the following:

Theorem ([SS03], Theorem 3.1.4)**.** *A collection of splits of $\{1,\dots,n\}$ is of the form $\Sigma(T)$ for a phylogenetic tree T on n taxa if and only if the splits are pairwise compatible.*

The irreducible boundary divisors in $\overline{M}_{0,n}$ are the codimension one strata given by dual graphs with a single internal edge; these are in bijection with splits $I \sqcup J = \{1,\dots,n\}$ such that I and J have cardinality at least two. Let us denote by $D_{I,J}$ the divisor corresponding to the split $I \sqcup J = \{1,\dots,n\}$, so that the generic point of $D_{I,J}$ corresponds to a two-component curve with a node separating the points indexed by I from those indexed by J. A pair of boundary divisors has nonempty intersection, $D_{I,J} \cap D_{K,L} \neq \emptyset$, if and only if at least one of the following containments holds [Kee92, pp.550–551]: $I \subset K$, $I \subset L$, $K \subset I$, $L \subset I$. Since $J = I^c$ and $L = K^c$, a pair of boundary divisors has nonempty intersection if and only if the corresponding splits are compatible.

Each stratum in $\overline{M}_{0,n}$ is the complete intersection of the boundary divisors containing it, and conversely any intersection of boundary divisors is a single stratum if it is nonempty. If a stratum has dual graph T, then the boundary divisors containing it are precisely the ones corresponding to splits in $\Sigma(T)$. Putting this together, we see that a collection of boundary divisors has nonempty intersection if and only if the corresponding set of splits is of the form $\Sigma(T)$ for some phylogenetic tree T, and by Buneman's result this is equivalent to the splits being pairwise compatible, which in turn is equivalent to the boundary divisors having pairwise nonempty intersection. □

References

[AFV11] Leonardo Aguirre, Giovanni Felder, and Alexander P. Veselov, *Gaudin subalgebras and stable rational curves*, Compos. Math. **147** (2011), no. 5, 1463–1478.

[BD86] Hans-Jürgen Bandelt and Andreas Dress, *Reconstructing the shape of a tree from observed dissimilarity data*, Adv. in Appl. Math. **7** (1986), no. 3, 309–343.

[BHV01] Louis Billera, Susan Holmes, and Karen Vogtmann, *Geometry of the space of phylogenetic trees*, Adv. in Appl. Math. **27** (2001), no. 4, 733–767.

[Bun71] P. Buneman, *The recovery of trees from measures of dissimilarity*, Mathematics in the Archaeological and Historical Sciencest, Edinburgh University Press, Edinburgh, 1971, pp. 387–395.

[Bun74] Peter Buneman, *A note on the metric properties of trees*, J. Combinatorial Theory Ser. B **17** (1974), 48–50.

[CT15] Ana-Maria Castravet and Jenia Tevelev, *$\overline{M}_{0,n}$ is not a Mori dream space*, Duke Math. J. **164** (2015), no. 8, 1641–1667.

[EHKR10] Pavel Etingof, André Henriques, Joel Kamnitzer, and Eric Rains, *The cohomology ring of the real locus of the moduli space of stable curves of genus 0 with marked points*, Ann. of Math. (2) **171** (2010), no. 2, 731–777.

[EJM76] G. Estabrook, C. Johnson, and F. McMorris, *A mathematical formulation for the analysis of cladistic character compatibility*, Math Bioscience **29** (1976).

[GM04] Alexander Goncharov and Yuri Manin, *Multiple ζ-motives and moduli spaces $\overline{M}_{0,n}$*, Compos. Math. **140** (2004), no. 1, 1–14.

[GM10] Angela Gibney and Diane Maclagan, *Equations for Chow and Hilbert quotients*, Algebra Number Theory **4** (2010), no. 7, 855–885.

[GRR72] Alexander Grothendieck, Michel Raynaud, and Dock Sang Rim, *Groupes de monodromie en géométrie algébrique. I*, Lecture Notes in Mathematics, Vol. 288, Springer-Verlag, 1972, Séminaire de Géométrie Algébrique du Bois-Marie 1967–1969 (SGA 7 I).

[Gus91] D. Gusfield, *Efficient algorithms for inferring evolutionary trees*, Networks **21** (1991), 19–28.

[HKT09] Paul Hacking, Sean Keel, and Jenia Tevelev, *Stable pair, tropical, and log canonical compactifications of moduli spaces of del Pezzo surfaces*, Invent. Math. **178** (2009), no. 1, 173–227.

[Kap93a] Mikhail Kapranov, *Chow quotients of Grassmannians. I*, I. M. Gelfand Seminar, Adv. Soviet Math., vol. 16, Amer. Math. Soc., Providence, RI, 1993, pp. 29–110.

[Kap93b] ———, *Veronese curves and Grothendieck-Knudsen moduli space $\overline{M}_{0,n}$*, J. Algebraic Geom. **2** (1993), no. 2, 239–262.

[Kap93c] Mikhail M. Kapranov, *The permutoassociahedron, Mac Lane's coherence theorem and asymptotic zones for the KZ equation*, J. Pure Appl. Algebra **85** (1993), no. 2, 119–142.

[Kee92] Sean Keel, *Intersection theory of moduli space of stable n-pointed curves of genus zero*, Trans. Amer. Math. Soc. **330** (1992), 545–574.

[KM13] Seán Keel and James McKernan, *Contractible extremal rays on $\overline{M}_{0,n}$*, Handbook of moduli. Vol. II, Adv. Lect. Math. (ALM), vol. 25, Int. Press, Somerville, MA, 2013, pp. 115–130.

[Knu83] Finn F. Knudsen, *The projectivity of the moduli space of stable curves. II. The stacks $M_{g,n}$*, Math. Scand. **52** (1983), no. 2, 161–199.

[Kol14] János Kollár, *Simple normal crossing varieties with prescribed dual complex*, Algebr. Geom. **1** (2014), no. 1, 57–68.

[KV07] Joachim Kock and Israel Vainsencher, *An invitation to quantum cohomology*, Progress in Mathematics, vol. 249, Birkhäuser Boston, Inc., Boston, MA, 2007, Kontsevich's formula for rational plane curves.

[LP11] Dan Levy and Lior Pachter, *The neighbor-net algorithm*, Adv. in Appl. Math. **47** (2011), no. 2, 240–258.

[Mea83] C. Meacham, *Theoretical and computational considerations of the compatibility of qualitative taxonomic characters*, Numerical taxonomy, NATO ASI Series, vol. G1, Springer-Verlag, 1983, pp. 304–314.

[MS13] Yu. I. Manin and M. N. Smirnov, *On the derived category of $\overline{M}_{0,n}$*, Izv. Ross. Akad. Nauk Ser. Mat. **77** (2013), no. 3, 93–108.

[Sch15] Luca Schaffler, *On the cone of effective 2-cycles on $\overline{M}_{0,7}$*, Eur. J. Math. **1** (2015), no. 4, 669–694.

[SS03] Charles Semple and Mike Steel, *Phylogenetics*, Oxford Lecture Series in Mathematics and its Applications, vol. 24, Oxford University Press, Oxford, 2003.

[SS04] David Speyer and Bernd Sturmfels, *The tropical Grassmannian*, Adv. Geom. **4** (2004), no. 3, 389–411.